



4845 Knightsbridge Blvd.
Suite 200
Columbus, OH 43214
Phone: (614) 451-4375
Fax: (614) 451-5284

Genetic Testing Summary

Enclosed are the genetic testing results for

CB 551

No amount of genetic testing can guarantee that a child will not be affected with a genetic condition. Genetic testing can inform you of the likelihood of passing on the genetic conditions that are tested for, but it cannot eliminate the risk of passing on any genetic condition.

The genetic conditions Cryobio tests for are inherited in an autosomal recessive manner. This means that the child would have to inherit a genetic mutation from both the sperm source and the egg source to be affected with the condition. When both the sperm source and the egg source have undergone genetic carrier screening and the test results are negative, the risk of a child being affected with the conditions tested for is significantly reduced, but it cannot be completely eliminated.

All recipients should discuss both their own risk for passing on genetic conditions and whether they would benefit from genetic counseling and testing with their health care provider. Before using a donor that is a carrier for a specific recessive genetic condition or conditions, we strongly recommend that the recipient (or egg source, if different) consider genetic counseling and testing to determine if they are a carrier for the same genetic condition or conditions as the donor.

Screening and testing have changed dramatically over the years, and so the screening and testing done on each donor may vary depending on the testing that was in place when he was actively in Cryobio's donor program. Earlier donors may not have had as extensive testing as later donors. Screening and testing may change again in the future, so please review the results each time before ordering as both the testing done and the results may change.



Patient Information

Name: Cb 551
Date of Birth: [REDACTED]
Sema4 ID: [REDACTED]
Client ID: [REDACTED]
Indication: Carrier Testing

Specimen Information

Specimen Type: Blood
Date Collected: [REDACTED]
Date Received: [REDACTED]
Final Report: [REDACTED]

Referring Provider

David Prescott, M.D.
Cryobiology, Inc.
4830-D Knightsbridge Blvd.
Columbus, OH, 43214
Fax: 614-451-5284

Expanded Carrier Screen (283)

Number of genes tested: 283

SUMMARY OF RESULTS AND RECOMMENDATIONS

⊕ Positive	⊖ Negative
<p>Carrier of Glycogen Storage Disease, Type V (AR) Associated gene(s): <i>PYGM</i> Variant(s) Detected: c.1094C>T, p.A365V, Likely Pathogenic, Heterozygous (one copy)</p> <p>Carrier of Joubert Syndrome 2 (AR) Associated gene(s): <i>TMEM216</i> Variant(s) Detected: c.253C>T, p.R85X, Pathogenic, Heterozygous (one copy)</p>	<p>Negative for all other genes tested To view a full list of genes and diseases tested please see Table 1 in this report</p>

AR=Autosomal recessive; XL=X-linked

Recommendations

- Testing the partner for the above positive disorder(s) and genetic counseling are recommended.
- Please note that for female carriers of X-linked diseases, follow-up testing of a male partner is not indicated.
- CGG repeat analysis of *FMR1* for fragile X syndrome is not performed on males as repeat expansion of premutation alleles is not expected in the male germline.
- Individuals of Asian, African, Hispanic and Mediterranean ancestry should also be screened for hemoglobinopathies by CBC and hemoglobin electrophoresis.
- Consideration of residual risk by ethnicity after a negative carrier screen is recommended for the other diseases on the panel, especially in the case of a positive family history for a specific disorder.

Interpretation of positive results

Glycogen Storage Disease, Type V (AR)

Results and Interpretation

A heterozygous (one copy) likely pathogenic missense variant, c.1094C>T, p.A365V, was detected in the *PYGM* gene (NM_005609.2). When this variant is present in trans with a pathogenic variant, it is considered to be causative for glycogen storage disease, type V. Therefore, this individual is expected to be at least a carrier for glycogen storage disease, type V. Heterozygous carriers are not expected to exhibit symptoms of this disease.

What is Glycogen Storage Disease, Type V?

Glycogen storage disease, type V (GSD5) is an autosomal recessive muscle disease caused by pathogenic variants in the gene *PYGM*. The clinical presentation is variable, although symptoms commonly begin in the first decade of life. Symptoms include exercise intolerance (pain when exercising) and muscle weakness, which may worsen in the patient's 40s or 50s. Most individuals are able to tolerate mild exercise and stay physically fit. Life expectancy is not affected. It is not currently possible to predict how severe the disease will be based on the inherited variants. GSD5 may affect people of any ethnicity, but it is more common among Sephardic Jewish individuals from Iran, Iraq and Syria.

Joubert Syndrome 2 (AR)

Results and Interpretation

A heterozygous (one copy) pathogenic premature stop codon, c.253C>T, p.R85X, was detected in the *TMEM216* gene (NM_001173990.2). When this variant is present in trans with a pathogenic variant, it is considered to be causative for Joubert syndrome 2. Therefore, this individual is expected to be at least a carrier for Joubert syndrome 2. Heterozygous carriers are not expected to exhibit symptoms of this disease.

What is Joubert Syndrome 2?

Joubert syndrome 2 is an autosomal recessive disorder caused by pathogenic variants in the gene *TMEM216*. While it has been identified in patients from different ethnicities, it is more prevalent in individuals of Ashkenazi Jewish descent, due to the presence of a founder mutation. Clinical features of Joubert syndrome include intellectual disability, brain malformations, ocular problems including uncontrollable eye movements and loss of vision, and kidney cysts leading to end-stage renal disease. Most patients with Joubert syndrome have a normal life expectancy. It is not currently possible to predict the severity of symptoms based on the genetic variants present.

Test description

This patient was tested for a panel of diseases using a combination of sequencing, targeted genotyping and copy number analysis. Please note that negative results reduce but do not eliminate the possibility that this individual is a carrier for one or more of the disorders tested. Please see Table 1 for a list of genes and diseases tested, and go.sema4.com/residualrisk for specific detection rates and residual risk by ethnicity. With individuals of mixed ethnicity, it is recommended to use the highest residual risk estimate. Only variants determined to be pathogenic or likely pathogenic are reported in this carrier screening test.



Xingwu Lu Ph.D., FACMG Assistant Laboratory Director

Laboratory Medical Consultant: George A. Diaz, M.D., Ph.D.

Genes and diseases tested

For specific detection rates and residual risk by ethnicity, please visit go.sema4.com/residualrisk

Table 1: List of genes and diseases tested with detailed results

Disease	Gene	Inheritance Pattern	Status	Detailed Summary
⊕ Positive				
Glycogen Storage Disease, Type V	PYGM	AR	Carrier	c.1094C>T, p.A365V, Likely Pathogenic, Heterozygous (one copy)
Joubert Syndrome 2	TMEM216	AR	Carrier	c.253C>T, p.R85X, Pathogenic, Heterozygous (one copy)
⊖ Negative				
3-Beta-Hydroxysteroid Dehydrogenase Type II Deficiency	HSD3B2	AR	Reduced Risk	
3-Methylcrotonyl-CoA Carboxylase Deficiency (MCCC1-Related)	MCCC1	AR	Reduced Risk	
3-Methylcrotonyl-CoA Carboxylase Deficiency (MCCC2-Related)	MCCC2	AR	Reduced Risk	
3-Methylglutaconic Aciduria, Type III	OPA3	AR	Reduced Risk	
3-Phosphoglycerate Dehydrogenase Deficiency	PHGDH	AR	Reduced Risk	
6-Pyruvoyl-Tetrahydropterin Synthase Deficiency	PTS	AR	Reduced Risk	
Abetalipoproteinemia	MTTP	AR	Reduced Risk	
Achromatopsia (CNGB3-related)	CNGB3	AR	Reduced Risk	
Acrodermatitis Enteropathica	SLC39A4	AR	Reduced Risk	
Acute Infantile Liver Failure	TRMU	AR	Reduced Risk	
Acyl-CoA Oxidase I Deficiency	ACOX1	AR	Reduced Risk	
Adenosine Deaminase Deficiency	ADA	AR	Reduced Risk	
Adrenoleukodystrophy, X-Linked	ABCD1	XL	Reduced Risk	
Aicardi-Goutieres Syndrome (SAMHD1-Related)	SAMHD1	AR	Reduced Risk	
Alpha-Mannosidosis	MAN2B1	AR	Reduced Risk	
Alpha-Thalassemia	HBA1/HBA2	AR	Reduced Risk	HBA1 Copy Number: 2 HBA2 Copy Number: 2 No pathogenic copy number variants detected HBA1/HBA2 Sequencing: Negative
Alpha-Thalassemia Mental Retardation Syndrome	ATRX	XL	Reduced Risk	
Alport Syndrome (COL4A3-Related)	COL4A3	AR	Reduced Risk	
Alport Syndrome (COL4A4-Related)	COL4A4	AR	Reduced Risk	
Alport Syndrome (COL4A5-Related)	COL4A5	XL	Reduced Risk	



Alstrom Syndrome	ALMS1	AR	Reduced Risk
Andermann Syndrome	SLC12A6	AR	Reduced Risk
Argininosuccinic Aciduria	ASL	AR	Reduced Risk
Aromatase Deficiency	CYP19A1	AR	Reduced Risk
Arthrogryposis, Mental Retardation, and Seizures	SLC35A3	AR	Reduced Risk
Asparagine Synthetase Deficiency	ASNS	AR	Reduced Risk
Aspartylglycosaminuria	AGA	AR	Reduced Risk
Ataxia With Isolated Vitamin E Deficiency	TTPA	AR	Reduced Risk
Ataxia-Telangiectasia	ATM	AR	Reduced Risk
Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay	SACS	AR	Reduced Risk
Bardet-Biedl Syndrome (BBS10-Related)	BBS10	AR	Reduced Risk
Bardet-Biedl Syndrome (BBS12-Related)	BBS12	AR	Reduced Risk
Bardet-Biedl Syndrome (BBS1-Related)	BBS1	AR	Reduced Risk
Bardet-Biedl Syndrome (BBS2-Related)	BBS2	AR	Reduced Risk
Bare Lymphocyte Syndrome, Type II	CIITA	AR	Reduced Risk
Barter Syndrome, Type 4A	BSND	AR	Reduced Risk
Bernard-Soulier Syndrome, Type A1	GP1BA	AR	Reduced Risk
Bernard-Soulier Syndrome, Type C	GP9	AR	Reduced Risk
Beta-Globin-Related Hemoglobinopathies	HBB	AR	Reduced Risk
Beta-Ketothiolase Deficiency	ACAT1	AR	Reduced Risk
Bilateral Frontoparietal Polymicrogyria	GPR56	AR	Reduced Risk
Biotinidase Deficiency	BTD	AR	Reduced Risk
Bloom Syndrome	BLM	AR	Reduced Risk
Canavan Disease	ASPA	AR	Reduced Risk
Carbamoylphosphate Synthetase I Deficiency	CPS1	AR	Reduced Risk
Carnitine Palmitoyltransferase IA Deficiency	CPT1A	AR	Reduced Risk
Carnitine Palmitoyltransferase II Deficiency	CPT2	AR	Reduced Risk
Carpenter Syndrome	RAB23	AR	Reduced Risk
Cartilage-Hair Hypoplasia	RMRP	AR	Reduced Risk
Cerebral Creatine Deficiency Syndrome 1	SLC6A8	XL	Reduced Risk
Cerebral Creatine Deficiency Syndrome 2	GAMT	AR	Reduced Risk
Cerebrotendinous Xanthomatosis	CYP27A1	AR	Reduced Risk
Charcot-Marie-Tooth Disease, Type 4D	NDRG1	AR	Reduced Risk
Charcot-Marie-Tooth Disease, Type 5 / Arts Syndrome	PRPS1	XL	Reduced Risk
Charcot-Marie-Tooth Disease, X-Linked	GJB1	XL	Reduced Risk



Choreoacanthocytosis	VPS13A	AR	Reduced Risk	
Choroideremia	CHM	XL	Reduced Risk	
Chronic Granulomatous Disease (CYBA-Related)	CYBA	AR	Reduced Risk	
Chronic Granulomatous Disease (CYBB-Related)	CYBB	XL	Reduced Risk	
Citrin Deficiency	SLC25A13	AR	Reduced Risk	
Citrullinemia, Type 1	ASS1	AR	Reduced Risk	
Cohen Syndrome	VPS13B	AR	Reduced Risk	
Combined Malonic and Methylmalonic Aciduria	ACSF3	AR	Reduced Risk	
Combined Oxidative Phosphorylation Deficiency 1	GFM1	AR	Reduced Risk	
Combined Oxidative Phosphorylation Deficiency 3	TSFM	AR	Reduced Risk	
Combined Pituitary Hormone Deficiency 2	PROP1	AR	Reduced Risk	
Combined Pituitary Hormone Deficiency 3	LHX3	AR	Reduced Risk	
Combined SAP Deficiency	PSAP	AR	Reduced Risk	
Congenital Adrenal Hyperplasia due to 17-Alpha-Hydroxylase Deficiency	CYP17A1	AR	Reduced Risk	
Congenital Adrenal Hyperplasia due to 21-Hydroxylase Deficiency	CYP21A2	AR	Reduced Risk	CYP21A2 copy number: 2 CYP21A2 sequencing: Negative
Congenital Amegakaryocytic Thrombocytopenia	MPL	AR	Reduced Risk	
Congenital Disorder of Glycosylation, Type Ia	PMM2	AR	Reduced Risk	
Congenital Disorder of Glycosylation, Type Ib	MPI	AR	Reduced Risk	
Congenital Disorder of Glycosylation, Type Ic	ALG6	AR	Reduced Risk	
Congenital Insensitivity to Pain with Anhidrosis	NTRK1	AR	Reduced Risk	
Congenital Myasthenic Syndrome (CHRNE-Related)	CHRNE	AR	Reduced Risk	
Congenital Myasthenic Syndrome (RAPSN-Related)	RAPSN	AR	Reduced Risk	
Congenital Neutropenia (HAX1-Related)	HAX1	AR	Reduced Risk	
Congenital Neutropenia (VPS45-Related)	VPS45	AR	Reduced Risk	
Corneal Dystrophy and Perceptive Deafness	SLC4A11	AR	Reduced Risk	
Corticosterone Methyloxidase Deficiency	CYP11B2	AR	Reduced Risk	
Cystic Fibrosis	CFTR	AR	Reduced Risk	
Cystinosis	CTNS	AR	Reduced Risk	
D-Bifunctional Protein Deficiency	HSD17B4	AR	Reduced Risk	
Deafness, Autosomal Recessive 77	LOXHD1	AR	Reduced Risk	
Duchenne Muscular Dystrophy / Becker Muscular Dystrophy	DMD	XL	Reduced Risk	
Dyskeratosis Congenita (RTEL1-Related)	RTEL1	AR	Reduced Risk	
Dystrophic Epidermolysis Bullosa	COL7A1	AR	Reduced Risk	
Ehlers-Danlos Syndrome, Type VIIC	ADAMTS2	AR	Reduced Risk	



Ellis-van Creveld Syndrome (EVC-Related)	<i>EVC</i>	AR	Reduced Risk	
Emery-Dreifuss Myopathy 1	<i>EMD</i>	XL	Reduced Risk	
Enhanced S-Cone Syndrome	<i>NR2E3</i>	AR	Reduced Risk	
Ethylmalonic Encephalopathy	<i>ETHE1</i>	AR	Reduced Risk	
Fabry Disease	<i>GLA</i>	XL	Reduced Risk	
Factor IX Deficiency	<i>F9</i>	XL	Reduced Risk	
Factor XI Deficiency	<i>F11</i>	AR	Reduced Risk	
Familial Autosomal Recessive Hypercholesterolemia	<i>LDLRAP1</i>	AR	Reduced Risk	
Familial Dysautonomia	<i>IKBKAP</i>	AR	Reduced Risk	
Familial Hypercholesterolemia	<i>LDLR</i>	AR	Reduced Risk	
Familial Hyperinsulinism (ABCC8-Related)	<i>ABCC8</i>	AR	Reduced Risk	
Familial Hyperinsulinism (KCNJ11-Related)	<i>KCNJ11</i>	AR	Reduced Risk	
Familial Mediterranean Fever	<i>MEFV</i>	AR	Reduced Risk	
Fanconi Anemia, Group A	<i>FANCA</i>	AR	Reduced Risk	
Fanconi Anemia, Group C	<i>FANCC</i>	AR	Reduced Risk	
Fanconi Anemia, Group G	<i>FANCG</i>	AR	Reduced Risk	
Fragile X Syndrome	<i>FMR1</i>	XL	Reduced Risk	<i>FMR1</i> CGG repeat sizes: Not Performed <i>FMR1</i> Sequencing: Negative Fragile X CGG triplet repeat expansion testing was not performed at this time, as the patient has either been previously tested or is a male.
Fumarase Deficiency	<i>FH</i>	AR	Reduced Risk	
GRACILE Syndrome and Other <i>BCS1L</i> -Related Disorders	<i>BCS1L</i>	AR	Reduced Risk	
Galactokinase Deficiency	<i>GALK1</i>	AR	Reduced Risk	
Galactosemia	<i>GALT</i>	AR	Reduced Risk	
Gaucher Disease	<i>GBA</i>	AR	Reduced Risk	
Gitelman Syndrome	<i>SLC12A3</i>	AR	Reduced Risk	
Glutaric Acidemia, Type I	<i>GCDH</i>	AR	Reduced Risk	
Glutaric Acidemia, Type IIa	<i>ETF A</i>	AR	Reduced Risk	
Glutaric Acidemia, Type IIc	<i>ETFDH</i>	AR	Reduced Risk	
Glycine Encephalopathy (AMT-Related)	<i>AMT</i>	AR	Reduced Risk	
Glycine Encephalopathy (GLDC-Related)	<i>GLDC</i>	AR	Reduced Risk	
Glycogen Storage Disease, Type II	<i>GAA</i>	AR	Reduced Risk	
Glycogen Storage Disease, Type III	<i>AGL</i>	AR	Reduced Risk	
Glycogen Storage Disease, Type IV / Adult Polyglucosan Body Disease	<i>GBE1</i>	AR	Reduced Risk	
Glycogen Storage Disease, Type Ia	<i>G6PC</i>	AR	Reduced Risk	
Glycogen Storage Disease, Type Ib	<i>SLC37A4</i>	AR	Reduced Risk	



Glycogen Storage Disease, Type VII	<i>PFKM</i>	AR	Reduced Risk
HMG-CoA Lyase Deficiency	<i>HMGCL</i>	AR	Reduced Risk
Hemochromatosis, Type 2A	<i>HFE2</i>	AR	Reduced Risk
Hemochromatosis, Type 3	<i>TFR2</i>	AR	Reduced Risk
Hereditary Fructose Intolerance	<i>ALDOB</i>	AR	Reduced Risk
Hereditary Spastic Paraparesis 49	<i>TECPR2</i>	AR	Reduced Risk
Hermansky-Pudlak Syndrome, Type 1	<i>HPS1</i>	AR	Reduced Risk
Hermansky-Pudlak Syndrome, Type 3	<i>HPS3</i>	AR	Reduced Risk
Holocarboxylase Synthetase Deficiency	<i>HLCS</i>	AR	Reduced Risk
Homocystinuria (CBS-Related)	<i>CBS</i>	AR	Reduced Risk
Homocystinuria due to <i>MTHFR</i> Deficiency	<i>MTHFR</i>	AR	Reduced Risk
Homocystinuria, cbIE Type	<i>MTRR</i>	AR	Reduced Risk
Hydroletharus Syndrome	<i>HYLS1</i>	AR	Reduced Risk
Hyperornithinemia-Hyperammonemia-Homocitrullinuria Syndrome	<i>SLC25A15</i>	AR	Reduced Risk
Hypohidrotic Ectodermal Dysplasia 1	<i>EDA</i>	XL	Reduced Risk
Hypophosphatasia	<i>ALPL</i>	AR	Reduced Risk
Inclusion Body Myopathy 2	<i>GNE</i>	AR	Reduced Risk
Infantile Cerebral and Cerebellar Atrophy	<i>MED17</i>	AR	Reduced Risk
Isovaleric Acidemia	<i>IVD</i>	AR	Reduced Risk
Joubert Syndrome 7 / Meckel Syndrome 5 / COACH Syndrome	<i>RPGRIP1L</i>	AR	Reduced Risk
Junctional Epidermolysis Bullosa (<i>LAMA3</i> -Related)	<i>LAMA3</i>	AR	Reduced Risk
Junctional Epidermolysis Bullosa (<i>LAMB3</i> -Related)	<i>LAMB3</i>	AR	Reduced Risk
Junctional Epidermolysis Bullosa (<i>LAMC2</i> -Related)	<i>LAMC2</i>	AR	Reduced Risk
Krabbe Disease	<i>GALC</i>	AR	Reduced Risk
Lamellar Ichthyosis, Type 1	<i>TGM1</i>	AR	Reduced Risk
Leber Congenital Amaurosis 10 and Other CEP290-Related Ciliopathies	<i>CEP290</i>	AR	Reduced Risk
Leber Congenital Amaurosis 13	<i>RDH12</i>	AR	Reduced Risk
Leber Congenital Amaurosis 2 / Retinitis Pigmentosa 20	<i>RPE65</i>	AR	Reduced Risk
Leber Congenital Amaurosis 5	<i>LCA5</i>	AR	Reduced Risk
Leber Congenital Amaurosis 8 / Retinitis Pigmentosa 12 / Pigmented Paravenous Chorioretinal Atrophy	<i>CRB1</i>	AR	Reduced Risk
Leigh Syndrome, French-Canadian Type	<i>LRPPRC</i>	AR	Reduced Risk
Lethal Congenital Contracture Syndrome 1 / Lethal Arthrogryposis with Anterior Horn Cell Disease	<i>GLE1</i>	AR	Reduced Risk
Leukoencephalopathy with Vanishing White Matter	<i>EIF2B5</i>	AR	Reduced Risk



Limb-Girdle Muscular Dystrophy, Type 2A	CAPN3	AR	Reduced Risk
Limb-Girdle Muscular Dystrophy, Type 2B	DYSF	AR	Reduced Risk
Limb-Girdle Muscular Dystrophy, Type 2C	SGCG	AR	Reduced Risk
Limb-Girdle Muscular Dystrophy, Type 2D	SGCA	AR	Reduced Risk
Limb-Girdle Muscular Dystrophy, Type 2E	SGCB	AR	Reduced Risk
Limb-Girdle Muscular Dystrophy, Type 2I	FKRP	AR	Reduced Risk
Lipoamide Dehydrogenase Deficiency	DLD	AR	Reduced Risk
Lipoid Adrenal Hyperplasia	STAR	AR	Reduced Risk
Lipoprotein Lipase Deficiency	LPL	AR	Reduced Risk
Long-Chain 3-Hydroxyacyl-CoA Dehydrogenase Deficiency	HADHA	AR	Reduced Risk
Lysinuric Protein Intolerance	SLC7A7	AR	Reduced Risk
Maple Syrup Urine Disease, Type 1a	BCKDHA	AR	Reduced Risk
Maple Syrup Urine Disease, Type 1b	BCKDHB	AR	Reduced Risk
Meckel 1 / Bardet-Biedl Syndrome 13	MKS1	AR	Reduced Risk
Medium Chain Acyl-CoA Dehydrogenase Deficiency	ACADM	AR	Reduced Risk
Megalencephalic Leukoencephalopathy with Subcortical Cysts	MLC1	AR	Reduced Risk
Menkes Disease	ATP7A	XL	Reduced Risk
Metachromatic Leukodystrophy	ARSA	AR	Reduced Risk
Methylmalonic Acidemia (MMAA-Related)	MMAA	AR	Reduced Risk
Methylmalonic Acidemia (MMAB-Related)	MMAB	AR	Reduced Risk
Methylmalonic Acidemia (MUT-Related)	MUT	AR	Reduced Risk
Methylmalonic Aciduria and Homocystinuria, Cobalamin C Type	MMACHC	AR	Reduced Risk
Methylmalonic Aciduria and Homocystinuria, Cobalamin D Type	MMADHC	AR	Reduced Risk
Microphthalmia / Anophthalmia	VSX2	AR	Reduced Risk
Mitochondrial Complex I Deficiency (ACAD9-Related)	ACAD9	AR	Reduced Risk
Mitochondrial Complex I Deficiency (NDUFA5-Related)	NDUFA5	AR	Reduced Risk
Mitochondrial Complex I Deficiency (NDUFS6-Related)	NDUFS6	AR	Reduced Risk
Mitochondrial DNA Depletion Syndrome 6 / Navajo Neurohepatopathy	MPV17	AR	Reduced Risk
Mitochondrial Myopathy and Sideroblastic Anemia 1	PUS1	AR	Reduced Risk
Mucopolidosis II / IIIA	GNPTAB	AR	Reduced Risk
Mucopolidosis III Gamma	GNPTG	AR	Reduced Risk
Mucopolidosis IV	MCOLN1	AR	Reduced Risk
Mucopolysaccharidosis Type I	IDUA	AR	Reduced Risk



Mucopolysaccharidosis Type II	IDS	XL	Reduced Risk
Mucopolysaccharidosis Type IIIA	SGSH	AR	Reduced Risk
Mucopolysaccharidosis Type IIIB	NAGLU	AR	Reduced Risk
Mucopolysaccharidosis Type IIIC	HGSNAT	AR	Reduced Risk
Mucopolysaccharidosis Type IIID	GNS	AR	Reduced Risk
Mucopolysaccharidosis Type IVb / GM1 Gangliosidosis	GLB1	AR	Reduced Risk
Mucopolysaccharidosis type IX	HYAL1	AR	Reduced Risk
Mucopolysaccharidosis type VI	ARSB	AR	Reduced Risk
Multiple Sulfatase Deficiency	SUMF1	AR	Reduced Risk
Muscle-Eye-Brain Disease and Other <i>POMGNT1</i> -Related Congenital Muscular Dystrophy-Dystroglycanopathies	<i>POMGNT1</i>	AR	Reduced Risk
Myoneurogastrointestinal Encephalopathy	TYMP	AR	Reduced Risk
Myotubular Myopathy 1	MTM1	XL	Reduced Risk
N-Acetylglutamate Synthase Deficiency	NAGS	AR	Reduced Risk
Nemaline Myopathy 2	NEB	AR	Reduced Risk
Nephrogenic Diabetes Insipidus, Type II	AQP2	AR	Reduced Risk
Nephrotic Syndrome (<i>NPHS1</i> -Related) / Congenital Finnish Nephrosis	<i>NPHS1</i>	AR	Reduced Risk
Nephrotic Syndrome (<i>NPHS2</i> -Related) / Steroid-Resistant Nephrotic Syndrome	<i>NPHS2</i>	AR	Reduced Risk
Neuronal Ceroid-Lipofuscinosis (<i>CLN3</i> -Related)	<i>CLN3</i>	AR	Reduced Risk
Neuronal Ceroid-Lipofuscinosis (<i>CLN5</i> -Related)	<i>CLN5</i>	AR	Reduced Risk
Neuronal Ceroid-Lipofuscinosis (<i>CLN6</i> -Related)	<i>CLN6</i>	AR	Reduced Risk
Neuronal Ceroid-Lipofuscinosis (<i>CLN8</i> -Related)	<i>CLN8</i>	AR	Reduced Risk
Neuronal Ceroid-Lipofuscinosis (<i>MFSD8</i> -Related)	<i>MFSD8</i>	AR	Reduced Risk
Neuronal Ceroid-Lipofuscinosis (<i>PPT1</i> -Related)	<i>PPT1</i>	AR	Reduced Risk
Neuronal Ceroid-Lipofuscinosis (<i>TPP1</i> -Related)	<i>TPP1</i>	AR	Reduced Risk
Niemann-Pick Disease (<i>SMPD1</i> -Related)	<i>SMPD1</i>	AR	Reduced Risk
Niemann-Pick Disease, Type C (<i>NPC1</i> -Related)	<i>NPC1</i>	AR	Reduced Risk
Niemann-Pick Disease, Type C (<i>NPC2</i> -Related)	<i>NPC2</i>	AR	Reduced Risk
Nijmegen Breakage Syndrome	<i>NBN</i>	AR	Reduced Risk
Non-Syndromic Hearing Loss (<i>GJB2</i> -Related)	<i>GJB2</i>	AR	Reduced Risk
Odonto-Onycho-Dermal Dysplasia / Schopf-Schulz-Passarge Syndrome	<i>WNT10A</i>	AR	Reduced Risk
Omenn Syndrome (<i>RAG2</i> -Related)	<i>RAG2</i>	AR	Reduced Risk
Omenn Syndrome / Severe Combined Immunodeficiency, Athabaskan-Type	<i>DCLRE1C</i>	AR	Reduced Risk
Ornithine Aminotransferase Deficiency	<i>OAT</i>	AR	Reduced Risk



Ornithine Transcarbonylase Deficiency	OTC	XL	Reduced Risk
Osteopetrosis 1	TCIRG1	AR	Reduced Risk
Pendred Syndrome	SLC26A4	AR	Reduced Risk
Phenylalanine Hydroxylase Deficiency	PAH	AR	Reduced Risk
Polycystic Kidney Disease, Autosomal Recessive	PKHD1	AR	Reduced Risk
Polyglandular Autoimmune Syndrome, Type 1	AIRE	AR	Reduced Risk
Pontocerebellar Hypoplasia, Type 1A	VRK1	AR	Reduced Risk
Pontocerebellar Hypoplasia, Type 6	RARS2	AR	Reduced Risk
Primary Carnitine Deficiency	SLC22A5	AR	Reduced Risk
Primary Ciliary Dyskinesia (DNAH5-Related)	DNAH5	AR	Reduced Risk
Primary Ciliary Dyskinesia (DNAI1-Related)	DNAI1	AR	Reduced Risk
Primary Ciliary Dyskinesia (DNAI2-Related)	DNAI2	AR	Reduced Risk
Primary Hyperoxaluria, Type 1	AGXT	AR	Reduced Risk
Primary Hyperoxaluria, Type 2	GRHPR	AR	Reduced Risk
Primary Hyperoxaluria, Type 3	HOGA1	AR	Reduced Risk
Progressive Cerebello-Cerebral Atrophy	SEPSECS	AR	Reduced Risk
Progressive Familial Intrahepatic Cholestasis, Type 2	ABCB11	AR	Reduced Risk
Propionic Acidemia (PCCA-Related)	PCCA	AR	Reduced Risk
Propionic Acidemia (PCCB-Related)	PCCB	AR	Reduced Risk
Pycnodysostosis	CTSK	AR	Reduced Risk
Pyruvate Dehydrogenase E1-Alpha Deficiency	PDHA1	XL	Reduced Risk
Pyruvate Dehydrogenase E1-Beta Deficiency	PDHB	AR	Reduced Risk
Renal Tubular Acidosis and Deafness	ATP6V1B1	AR	Reduced Risk
Retinitis Pigmentosa 25	EYS	AR	Reduced Risk
Retinitis Pigmentosa 26	CERKL	AR	Reduced Risk
Retinitis Pigmentosa 28	FAM161A	AR	Reduced Risk
Retinitis Pigmentosa 59	DHDDS	AR	Reduced Risk
Rhizomelic Chondrodysplasia Punctata, Type 1	PEX7	AR	Reduced Risk
Rhizomelic Chondrodysplasia Punctata, Type 3	AGPS	AR	Reduced Risk
Roberts Syndrome	ESCO2	AR	Reduced Risk
Salla Disease	SLC17A5	AR	Reduced Risk
Sandhoff Disease	HEXB	AR	Reduced Risk
Schimke Immunoosseous Dysplasia	SMARCAL1	AR	Reduced Risk
Segawa Syndrome	TH	AR	Reduced Risk
Sjogren-Larsson Syndrome	ALDH3A2	AR	Reduced Risk
Smith-Lemli-Opitz Syndrome	DHCR7	AR	Reduced Risk



Spinal Muscular Atrophy	SMN1	AR	Reduced Risk	SMN1 copy number: 2 SMN2 copy number: 1 c. *3+80T>G: Negative
Spondylothoracic Dysostosis	MESP2	AR	Reduced Risk	
Steel Syndrome	COL27A1	AR	Reduced Risk	
Stuve-Wiedemann Syndrome	LIFR	AR	Reduced Risk	
Sulfate Transporter-Related Osteochondrodysplasia	SLC26A2	AR	Reduced Risk	
Tay-Sachs Disease	HEXA	AR	Reduced Risk	Tay-Sachs disease enzyme: Non-carrier White blood cells: Non-carrier <ul style="list-style-type: none"> Hex A%: 67.1% (Non-carrier: 55.0 - 72.0%; Carrier: <50%) Total hexosaminidase activity: 1301 nmol/hr/mg Plasma: Non-carrier <ul style="list-style-type: none"> Hex A%: 70.1 (Non-carrier: 58.0 - 72.0%; Carrier: <54%) Total hexosaminidase activity: 538 nmol/hr/ml HEXA Sequencing: Negative
Tyrosinemia, Type I	FAH	AR	Reduced Risk	
Usher Syndrome, Type IB	MYO7A	AR	Reduced Risk	
Usher Syndrome, Type IC	USH1C	AR	Reduced Risk	
Usher Syndrome, Type ID	CDH23	AR	Reduced Risk	
Usher Syndrome, Type IF	PCDH15	AR	Reduced Risk	
Usher Syndrome, Type IIA	USH2A	AR	Reduced Risk	
Usher Syndrome, Type III	CLRN1	AR	Reduced Risk	
Very Long Chain Acyl-CoA Dehydrogenase Deficiency	ACADVL	AR	Reduced Risk	
Walker-Warburg Syndrome and Other FKTN-Related Dystrophies	FKTN	AR	Reduced Risk	
Wilson Disease	ATP7B	AR	Reduced Risk	
Wolman Disease / Cholesteryl Ester Storage Disease	LIPA	AR	Reduced Risk	
X-Linked Juvenile Retinoschisis	RS1	XL	Reduced Risk	
X-Linked Severe Combined Immunodeficiency	IL2RG	XL	Reduced Risk	
Zellweger Syndrome Spectrum (PEX10-Related)	PEX10	AR	Reduced Risk	
Zellweger Syndrome Spectrum (PEX1-Related)	PEX1	AR	Reduced Risk	
Zellweger Syndrome Spectrum (PEX2-Related)	PEX2	AR	Reduced Risk	
Zellweger Syndrome Spectrum (PEX6-Related)	PEX6	AR	Reduced Risk	

AR=Autosomal recessive; XL=X-linked

Test methods and comments

Genomic DNA isolated from this patient was analyzed by one or more of the following methodologies, as applicable:

Fragile X CGG Repeat Analysis (Analytical Detection Rate >99%)

PCR amplification using Asuragen, Inc. AmplideX® *FMR1* PCR reagents followed by capillary electrophoresis for allele sizing was performed. Samples positive for *FMR1* CGG repeats in the premutation and full mutation size range were further analyzed by Southern blot analysis to assess the size and methylation status of the *FMR1* CGG repeat.

Genotyping (Analytical Detection Rate >99%)

Multiplex PCR amplification and allele specific primer extension analyses using the MassARRAY® System were used to identify variants that are complex in nature or are present in low copy repeats. Rare sequence variants may interfere with assay performance.

Multiplex Ligation-Dependent Probe Amplification (MLPA) (Analytical Detection Rate >99%)

MLPA® probe sets and reagents from MRC-Holland were used for copy number analysis of specific targets versus known control samples. False positive or negative results may occur due to rare sequence variants in target regions detected by MLPA probes. Analytical sensitivity and specificity of the MLPA method are both 99%.

For alpha thalassemia, the copy numbers of the *HBA1* and *HBA2* genes were analyzed. Alpha-globin gene deletions, triplications, and the Constant Spring (CS) mutation are assessed. This test is expected to detect approximately 90% of all alpha-thalassemia mutations, varying by ethnicity. Carriers of alpha-thalassemia with three or more *HBA* copies on one chromosome, and one or no copies on the other chromosome, may not be detected. With the exception of triplications, other benign alpha-globin gene polymorphisms will not be reported. Analyses of *HBA1* and *HBA2* are performed in association with long-range PCR of the coding regions followed by short-read sequencing.

For Duchenne muscular dystrophy, the copy numbers of all *DMD* exons were analyzed. Potentially pathogenic single exon deletions and duplications are confirmed by a second method. Analysis of *DMD* is performed in association with sequencing of the coding regions.

For congenital adrenal hyperplasia, the copy number of the *CYP21A2* gene was analyzed. This analysis can detect large deletions due to unequal meiotic crossing-over between *CYP21A2* and the pseudogene *CYP21A1P*. These 30-kb deletions make up approximately 20% of *CYP21A2* pathogenic alleles. This test may also identify certain point mutations in *CYP21A2* caused by gene conversion events between *CYP21A2* and *CYP21A1P*. Some carriers may not be identified by dosage sensitive methods as this testing cannot detect individuals with two copies (duplication) of the *CYP21A2* gene on one chromosome and loss of *CYP21A2* (deletion) on the other chromosome. Analysis of *CYP21A2* is performed in association with long-range PCR of the coding regions followed by short-read sequencing.

For spinal muscular atrophy (SMA), the copy numbers of the *SMN1* and *SMN2* genes were analyzed. The individual dosage of exons 7 and 8 as well as the combined dosage of exons 1, 4, 6 and 8 of *SMN1* and *SMN2* were assessed. Copy number gains and losses can be detected with this assay. Depending on ethnicity, 6 - 29% of carriers will not be identified by dosage sensitive methods as this testing cannot detect individuals with two copies (duplication) of the *SMN1* gene on one chromosome and loss of *SMN1* (deletion) on the other chromosome (silent 2+0 carrier) or individuals that carry an intragenic mutation in *SMN1*. Please also note that 2% of individuals with SMA have an *SMN1* mutation that occurred *de novo*. Typically in these cases, only one parent is an SMA carrier.

The presence of the c.*3+80T>G (chr5:70,247,901T>G) variant allele in an individual with Ashkenazi Jewish or Asian ancestry is typically indicative of a duplication of *SMN1*. When present in an Ashkenazi Jewish or Asian individual with two copies of *SMN1*, c.*3+80T>G is likely indicative of a silent (2+0) carrier. In individuals with two copies of *SMN1* with African American, Hispanic or Caucasian ancestry, the presence or absence of c.*3+80T>G significantly increases or decreases, respectively, the likelihood of being a silent 2+0 carrier.

Pathogenic or likely pathogenic sequence variants in exon 7 may be detected during testing for the c.*3+80T>G variant allele; these will be reported if confirmed to be located in *SMN1* using locus-specific Sanger primers

MLPA for Gaucher disease (*GBA*), cystic fibrosis (*CFTR*), and non-syndromic hearing loss (*GJB2/GJB6*) will only be performed if indicated for confirmation of detected CNVs. If *GBA* analysis was performed, the copy numbers of exons 1, 3, 4, and 6 - 10 of the *GBA* gene (of 11 exons total) were analyzed. If *CFTR* analysis was performed, the copy numbers of all 27 *CFTR* exons were analyzed. If *GJB2/GJB6* analysis was performed, the copy number of the two *GJB2* exons were analyzed, as well as the presence or absence of the two upstream deletions of the *GJB2* regulatory region, del(*GJB6*-D13S1830) and del(*GJB6*-D13S1854).

Next Generation Sequencing (NGS) (Analytical Detection Rate >95%)

NGS was performed on a panel of genes for the purpose of identifying pathogenic or likely pathogenic variants.

Agilent SureSelect™ QXT technology was used with a custom capture library to target the exonic regions and intron/exon splice junctions of the relevant genes, as well as a number of UTR, intronic or promoter regions that contain previously reported mutations. Samples were pooled and sequenced on the Illumina HiSeq 2500 platform in the Rapid Run mode or the Illumina NovaSeq platform in the Xp workflow, using 100 bp paired-end reads. The sequencing data was analyzed using a custom bioinformatics algorithm designed and validated in house.

The coding exons and splice junctions of the known protein-coding RefSeq genes were assessed for the average depth of coverage (minimum of 20X) and data quality threshold values. Most exons not meeting a minimum of >20X read depth across the exon are further analyzed by Sanger sequencing. Please note that several genomic regions present difficulties in mapping or obtaining read depth >20X. The exons contained within these regions are noted within Table 1 (as "Exceptions") and will not be reflexed to Sanger sequencing if the mapping



quality or coverage is poor. Any variants identified during testing in these regions are confirmed by a second method and reported if determined to be pathogenic or likely pathogenic. However, as there is a possibility of false negative results within these regions, detection rates and residual risks for these genes have been calculated with the presumption that variants in these exons will not be detected, unless included in the MassARRAY® genotyping platform.

This test will detect variants within the exons and the intron-exon boundaries of the target regions. Variants outside these regions may not be detected, including, but not limited to, UTRs, promoters, and deep intronic areas, or regions that fall into the Exceptions mentioned above. This technology may not detect all small insertion/deletions and is not diagnostic for repeat expansions and structural genomic variation. In addition, a mutation(s) in a gene not included on the panel could be present in this patient.

Variant interpretation and classification was performed based on the American College of Medical Genetics Standards and Guidelines for the Interpretation of Sequence Variants (Richards et al, 2015). All potentially pathogenic variants may be confirmed by either a specific genotyping assay or Sanger sequencing, if indicated. Any benign variants, likely benign variants or variants of uncertain significance identified during this analysis will not be reported.

Copy Number Variant Analysis (Analytical Detection Rate >95%)

Large duplications and deletions were called from the relative read depths on an exon-by-exon basis using a custom exome hidden Markov model (XHMM) algorithm. Deletions or duplications determined to be pathogenic or likely pathogenic were confirmed by either a custom arrayCGH platform, quantitative PCR, or MLPA (depending on CNV size and gene content). While this algorithm is designed to pick up deletions and duplications of 2 or more exons in length, potentially pathogenic single-exon CNVs will be confirmed and reported, if detected.

Exon Array (Confirmation method) (Accuracy >99%)

The customized oligonucleotide microarray (Oxford Gene Technology) is a highly-targeted exon-focused array capable of detecting medically relevant microdeletions and microduplications at a much higher resolution than traditional aCGH methods. Each array matrix has approximately 180,000 60-mer oligonucleotide probes that cover the entire genome. This platform is designed based on human genome NCBI Build 37 (hg19) and the CGH probes are enriched to target the exonic regions of the genes in this panel.

Quantitative PCR (Confirmation method) (Accuracy >99%)

The relative quantification PCR is utilized on a Roche Universal Library Probe (UPL) system, which relates the PCR signal of the target region in one group to another. To test for genomic imbalances, both sample DNA and reference DNA is amplified with primer/probe sets that specific to the target region and a control region with known genomic copy number. Relative genomic copy numbers are calculated based on the standard $\Delta\Delta C_t$ formula.

Long-Range PCR (Analytical Detection Rate >99%)

Long-range PCR was performed to generate locus-specific amplicons for *CYP21A2*, *HBA1* and *HBA2* and *GBA*. The PCR products were then prepared for short-read NGS sequencing and sequenced. Sequenced reads were mapped back to the original genomic locus and run through the bioinformatics pipeline. If indicated, copy number from MLPA was correlated with the sequencing output to analyze the results. For *CYP21A2*, a certain percentage of healthy individuals carry a duplication of the *CYP21A2* gene, which has no clinical consequences. In cases where two copies of a gene are located on the same chromosome in tandem, only the second copy will be amplified and assessed for potentially pathogenic variants, due to size limitations of the PCR reaction. However, because these alleles contain at least two copies of the *CYP21A2* gene in tandem, it is expected that this patient has at least one functional gene in the tandem allele and this patient is therefore less likely to be a carrier. When an individual carries both a duplication allele and a pathogenic variant, or multiple pathogenic variants, the current analysis may not be able to determine the phase (cis/trans configuration) of the *CYP21A2* alleles identified. Family studies may be required in certain scenarios where phasing is required to determine the carrier status.

Residual Risk Calculations

Carrier frequencies and detection rates for each ethnicity were calculated through the combination of internal curations of >28,000 variants and genomic frequency data from >138,000 individuals across seven ethnic groups in the gnomAD database. Additional variants in HGMD and novel deleterious variants were also incorporated into the calculation. Residual risk values are calculated using a Bayesian analysis combining the *a priori* risk of being a pathogenic mutation carrier (carrier frequency) and the detection rate. They are provided only as a guide for assessing approximate risk given a negative result, and values will vary based on the exact ethnic background of an individual. This report does not represent medical advice but should be interpreted by a genetic counselor, medical geneticist or physician skilled in genetic result interpretation and the relevant medical literature.

Sanger Sequencing (Confirmation method) (Accuracy >99%)

Sanger sequencing, as indicated, was performed using BigDye Terminator chemistry with the ABI 3730 DNA analyzer with target specific amplicons. It also may be used to supplement specific guaranteed target regions that fail NGS sequencing due to poor quality or low depth of coverage (<20 reads) or as a confirmatory method for NGS positive results. False negative results may occur if rare variants interfere with amplification or annealing.



Tay-Sachs Disease (TSD) Enzyme Analysis (Analytical Detection Rate \geq 98%)

Hexosaminidase activity and Hex A% activity were measured by a standard heat-inactivation, fluorometric method using artificial 4-MU- β -N-acetyl glucosaminide (4-MUG) substrate. This assay is highly sensitive and accurate in detecting Tay-Sachs carriers and individuals affected with TSD. Normal ranges of Hex A% activity are 55.0-72.0 for white blood cells and 58.0-72.0 for plasma. It is estimated that less than 0.5% of Tay-Sachs carriers have non-carrier levels of percent Hex A activity, and therefore may not be identified by this assay. In addition, this assay may detect individuals that are carriers of or are affected with Sandhoff disease. False positive results may occur if benign variants, such as pseudodeficiency alleles, interfere with the enzymatic assay. False negative results may occur if both *HEXA* and *HEXB* pathogenic or pseudodeficiency variants are present in the same individual.

Please note these tests were developed and their performance characteristics were determined by Mount Sinai Genomics, Inc. They have not been cleared or approved by the FDA. These analyses generally provide highly accurate information regarding the patient's carrier or affected status. Despite this high level of accuracy, it should be kept in mind that there are many potential sources of diagnostic error, including misidentification of samples, polymorphisms, or other rare genetic variants that interfere with analysis. Families should understand that rare diagnostic errors may occur for these reasons.

SELECTED REFERENCES

Carrier Screening

Grody W et al. ACMG position statement on prenatal/preconception expanded carrier screening. *Genet Med*. 2013 15:482-3.

Fragile X syndrome:

Chen L et al. An information-rich CGG repeat primed PCR that detects the full range of Fragile X expanded alleles and minimizes the need for Southern blot analysis. *J Mol Diag* 2010 12:589-600.

Spinal Muscular Atrophy:

Luo M et al. An Ashkenazi Jewish SMN1 haplotype specific to duplication alleles improves pan-ethnic carrier screening for spinal muscular atrophy. *Genet Med*. 2014 16:149-56.

Ashkenazi Jewish Disorders:

Scott SA et al. Experience with carrier screening and prenatal diagnosis for sixteen Ashkenazi Jewish Genetic Diseases. *Hum. Mutat*. 2010 31:1-11.


Duchenne Muscular Dystrophy:

Flanigan KM et al. Mutational spectrum of DMD mutations in dystrophinopathy patients: application of modern diagnostic techniques to a large cohort. *Hum Mutat*. 2009 30:1657-66.

Variant Classification:

Richards S et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015 May;17(5):405-24

Additional disease-specific references available upon request.



Cystic Fibrosis Mutation Analysis

Patient Name: CB, 551
Referring Physician: David Prescott, MD
Specimen #: [REDACTED]
Patient ID: [REDACTED]

Client #: [REDACTED]
Case #: [REDACTED]

[REDACTED]
Cryobiology, Inc.
4830-D Knightsbridge Boulevard
Columbus, OH 43214

DOB: [REDACTED] **Date Collected:** [REDACTED]
Sex: M **Date Received:** [REDACTED]
SSN: [REDACTED] **LAB ID:**
Hospital ID:
Specimen Type: BLDPER

Ethnicity: Caucasian

Indication: Carrier Test / Gamete donor

RESULTS: Negative for the 97 mutations analyzed

INTERPRETATION:

This individual is negative for the mutations analyzed. This result reduces but does not eliminate the risk to be a CF carrier. See Comments for ethnic-specific risk reductions based on a negative family history.

COMMENTS:

Mutations Detection Rates among Ethnic Groups		Detection Rates are based on mutation frequencies in patients affected with cystic fibrosis. Among individuals with an atypical or mild presentation (e.g. congenital absence of the vas deferens, pancreatitis) detection rates may vary from those provided here.	
Ethnicity	Carrier risk reduction when no family history	Detection rate	References
African American	1/61 to 1/316	81%	ACOG Committee Opinion 486 PMID: 21422883; Heim PMID: 11388756
Ashkenazi Jewish	1/24 to 1/767	97%	ACOG Committee Opinion 486 PMID: 21422883
Asian American	1/94 to <1/183	49-55%	ACOG Committee Opinion 486 PMID: 21422883; Watson PMID: 1384328
Caucasian	1/25 to 1/343	93%	ACOG Committee Opinion 486 PMID: 21422883; Heim PMID: 11388756; Palomaki PMID: 11882786
Hispanic	1/58 to 1/260	78%	ACOG Committee Opinion 486 PMID: 21422883; Heim PMID: 11388756; California Database: (http://www.cdph.ca.gov/programs/GDSP/Documents/CFTableCurrent.pdf)
Jewish, non-Ashkenazi		Varies by country of origin	Orgad PMID: 11336401; Kerem PMID:10464623
Mixed or Other		Not Provided	For counseling, consider using the ethnic background with the most conservative risk estimates.

This interpretation is based on the clinical and family relationship information provided and the current understanding of the molecular genetics of this condition.

METHOD / LIMITATIONS:

CFTR gene regions are amplified enzymatically. The 97 CF mutations are tested by multiplex allele-specific primer extension, bead array hybridization, and fluorescence detection. The test discriminates between p.F508del and three polymorphisms (p.I506V, p.I507V and p.F508C). Numbering and nomenclature follow Human Genome Variation Society recommendations. Mutations and their legacy names are listed at www.integratedgenetics.com/CFplus. The DNA reference sequence is NG_016465.1. False positive or negative results may occur for reasons that include genetic variants, blood transfusions, bone marrow transplantation, erroneous representation of family relationships, or maternal contamination of a fetal sample.

Integrated Genetics is a business unit of Esoterix Genetic Laboratories, LLC, a wholly-owned subsidiary of Laboratory Corporation of America Holdings.

Electronically Signed By: Zhaoqing Zhou, Ph.D., FACMG, on [REDACTED]

TO:Cryobiology, Inc. ATTN:Cryobiology, Inc.

MUTATIONS ANALYZED

c.54-5940_273+10250del21kb	c.1973_1985del13insAGAAA	p.R117H	p.R553*
c.262_263delTT	c.1976delA	p.Y122*	p.A559T
c.273+1G>A	c.2012delT	p.G178R	p.R560T
c.273+3A>C	c.2051_2052delAAinsG	p.L206W	p.P574H
c.274-1G>A	c.2052delA	p.F312del	p.R709*
c.313delA	c.2052dupA	p.G330*	p.K710*
c.325_327delTATinsG	c.2175dupA	p.R334W	p.R764*
c.442delA	c.2657+5G>A	p.T338I	p.Q890*
c.489+1G>T	c.2737_2738insG	p.R347H	p.R1066C
c.531delT	c.2988G>A	p.R347P	p.W1089*
c.579+1G>T	c.2988+1G>A	p.R352Q	p.Y1092*
c.579+5G>A	c.3039delC	p.[Q359K;T360K]	p.M1101K
c.580-1G>T	c.3067_3072delATAGTG	p.S364P	p.D1152H
c.803delA	c.3528delC	p.A455E	p.R1158*
c.805_806delAT	c.3536_3539delCCAA	p.G480C	p.R1162*
c.948delT	c.3659delC	p.Q493*	p.S1196*
c.1155_1156dupTA	c.3717+12191C>T	p.I507del	p.W1204*
c.1545_1546delTA	c.3744delA	p.F508del	p.Q1238*
c.1585-1G>A	c.3773dupT	p.V520F	p.S1251N
c.1680-1G>A	c.3889dupT	p.C524*	p.S1255*
c.1766+1G>A	p.E60*	p.G542*	p.W1282*
c.1766+5G>T	p.R75*	p.S549N	p.N1303K
c.1820_1903del84	p.G85E	p.S549R	
c.1911delG	p.E92*	p.G551D	
c.1923_1931del9insA	p.R117C	p.Q552*	

This test was developed and its performance characteristics determined by Esoterix Genetic Laboratories, LLC. It has not been cleared or approved by the Food and Drug Administration. The FDA has determined that such clearance or approval is not necessary.

Patient	Sample	Referring Doctor
Patient Name: Cb 553 Date of Birth: [REDACTED] Reference #: [REDACTED] Indication: Encounter of male for testing for genetic disease carrier status for procreative management (Z31.440) Test Type: Chromosome Analysis - Peripheral Blood	Specimen Type: Peripheral Blood Lab #: [REDACTED] Date Collected: [REDACTED] Date Received: [REDACTED] Final Report: [REDACTED]	David Prescott, M.D. Cryobiology, Inc. 4830-D Knightsbridge Blvd. Columbus, OH 43214 Fax: 614-451-5284

CYTOGENETIC ANALYSIS

Results

Staining: G-bands by trypsin using Giemsa (GTG)	Chromosome count: 46	Cells captured: 5
Band level: 550	Cells analyzed: 20	Cells karyotyped: 3
		Cultures examined: 2


Karyotype: **46,XY**

Interpretation

Cytogenetic analysis revealed the presence of a **normal male** karyotype in peripheral blood lymphocytes. This analysis does not show any evidence of a clinically significant numerical or structural chromosome abnormality.

The standard procedures used in this analysis do not routinely detect microdeletions, small rearrangements or low level mosaicism.

This case has been reviewed and electronically signed by Ram Singh, PhD, Assistant Laboratory Director
 Laboratory Medical Consultant: Bryn Webb, M.D.



SMN1 Copy Number Analysis

Patient Name: 551 CB

DOB: [REDACTED]

Age: [REDACTED]

SSN #: [REDACTED]

Gender: Male

[REDACTED]
Cryobiology, Inc.
4830-D Knightsbridge Boulevard
Columbus, OH 43214

Specimen #: [REDACTED]

Case #: [REDACTED]

Patient ID #: [REDACTED]

Date Collected: [REDACTED]

Date Received: [REDACTED]

Referring Physician: David Prescott

Client Lab ID #:

Genetic Counselor:

Hospital ID #:

Specimen Type: Peripheral Blood

Specimen ID #:

Specimen(s) Received: 1 - Lavender 7 ml round bottom tube(s)

Clinical Data: Carrier Test/Gamete donor

Ethnicity: Caucasian

RESULTS: SMN1 copy number: 2 (Reduced Carrier Risk)

INTERPRETATION:

This individual has an SMN1 copy number of two. This result reduces but does not eliminate the risk to be a carrier of SMA. Ethnic specific risk reductions based on a negative family history and an SMN1 copy number of two are provided in the Comments section of this report.

COMMENT:

Spinal muscular atrophy (SMA) is an autosomal recessive disease of variable age of onset and severity caused by mutations (most often deletions or gene conversions) in the survival motor neuron (SMN1) gene. Molecular testing assesses the number of copies of the SMN1 gene. Individuals with one copy of the SMN1 gene are predicted to be carriers of SMA. Individuals with two or more copies have a reduced risk to be carriers. (Affected individuals have 0 copies of the SMN1 gene.)

This copy number analysis cannot detect individuals who are carriers of SMA as a result of either 2 (or very rarely 3) copies of the SMN1 gene on one chromosome and the absence of the SMN1 gene on the other chromosome or small intragenic mutations within the SMN1 gene. This analysis also will not detect germline mosaicism or mutations in genes other than SMN1. Additionally, de novo mutations have been reported in approximately 2% of SMA patients.

Carrier Frequency and Risk Reductions for Individuals with No Family History of SMA

Ethnicity	Detection Rate ¹	Prior Carrier Risk ¹	Reduced Carrier Risk for 2 copy result	Reduced Carrier Risk for 3 copy result
Caucasian	94.8%	1:47	1:834	1:5,600
Ashkenazi Jewish	90.5%	1:67	1:611	1:5,400
Asian	93.3%	1:59	1:806	1:5,600
Hispanic	90.0%	1:68	1:579	1:5,400
African American	70.5%	1:72	1:130	1:4,200
Asian Indian	90.2%	1:52	1:443	1:5,400
Mixed or Other Ethnic Background	For counseling purposes, consider using the ethnic background with the most conservative risk estimates.			

METHOD/LIMITATIONS: Specimen DNA is isolated and amplified by real-time polymerase chain reaction (PCR) for exon 7 of the SMN1 gene and the internal standard reference genes. A mathematical algorithm is used to calculate and report SMN1 copy numbers of 0, 1, 2 and 3. Based upon this analysis, an upper limit of 3 represents the highest degree of accuracy in reporting SMN1 copy number with statistical confidence. Sequencing of the primer and probe binding sites is performed on all fetal samples and samples with one copy of SMN1 by real-time PCR to rule out the presence of sequence variants which could interfere with analysis and interpretation. False positive or negative results may occur for reasons that include genetic variants, blood transfusions, bone marrow transplantation, erroneous representation of family relationships or contamination of a fetal sample with maternal cells.

REFERENCES:

1. Sugarman EA, Nagan N, Zhu H, et al. Pan-ethnic carrier screening and prenatal diagnosis for spinal muscular atrophy: clinical laboratory analysis of >72,400 specimens. Eur J Hum Genet 2012; 20:27-32.
2. Prior TW, et al. Technical standards and guidelines for spinal muscular atrophy testing. Genet Med 2011; 13(7): 686-694.

The test was developed and its performance characteristics have been determined by Esoterix Genetic Laboratories, LLC. The laboratory is regulated under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high complexity clinical testing. This test must be used in conjunction with clinical assessment, when available. Integrated Genetics is a business unit of Esoterix Genetic Laboratories, LLC, a wholly-owned subsidiary of Laboratory Corporation of America Holdings.

Electronically Signed by: Zhaoqing Zhou, Ph.D., FACMG, on [REDACTED]

Reported by: /